

**LETTER TO THE EDITOR**

Reply to “racial and gender disparities among patients with Takotsubo syndrome”

To the Editor,

We read with great interest the Letter to the Editor titled “Racial and Gender Disparities among Patients with Takotsubo Syndrome” by Khalid et al¹ regarding our recent publication.² Their excellent comments and detailed assessment highlights the low prevalence of diabetes mellitus in patients with Takotsubo syndrome (TTS) compared to general population. This is in contrast with relatively high prevalence of many other cardiovascular risk factors in TTS patients. This so called “diabetes paradox” has been previously explained in TTS patients and is the target of many active investigations. As highlighted in the Letter to the Editor, the prevalence of diabetes mellitus in our patient population is very close to the results of prior meta-analyses of multiple small studies of patients with TTS.

We would also want to thank the authors for highlighting and discussing the difference in the rate of in-hospital mortality in male patients as compared to females. We agree with the proposed theory that attributes higher mortality in males to higher incidence of critical illnesses among them with resultant higher stress and elevated catecholamines. This is evident in our patient population as well but other investigators have suggested different explanations as to why male patients with TTS are more prone to morbidity and mortality. For example, in younger patient populations, TTS has been commonly reported in the setting of stimulant drug use or drug or alcohol withdrawal. Higher prevalence of these exposures in male population with TTS has been previously suggested as a possible theoretical explanation for higher mortality in them³ but as Khalid and colleagues suggest, we agree that more prospective trials based on multicenter, international registries are needed with special focus on risk factors and predictors of worse outcome and complications in TTS. These studies not only will help the clinical aspects of patient care, but also might provide glimpses into the elusive pathophysiology of TTS.

ACKNOWLEDGMENTS

None of the authors has any disclosures relevant to the content of the manuscript and has no potential conflict of interest to declare. No study specific funding was used to support this work. The authors are solely responsible for the study design, conduct and analyses, drafting and editing of the manuscript, and its final contents. All authors had access to the data and a role in writing the manuscript.

ORCID

Alejandro Lemor  <https://orcid.org/0000-0002-4649-8479>

Alejandro Lemor^{1,2} 

Seyed H. H. Dehkordi³

¹Division of Cardiology, Henry Ford Health Care System, Detroit, Michigan

²Centro de Investigación en Epidemiología Clínica y Medicina Basada en Evidencias, Facultad de Medicina Humana, Universidad de San Martín de Porres, Lima, Peru

³Division of Cardiology, University of Kansas Medical Center, Kansas City, Kansas

Correspondence

Alejandro Lemor, MD, MS, Division of Cardiology, Henry Ford Hospital, K-14, 2799 West Grand Blvd., Detroit, MI 48202.

Email: alejandrolemor@hotmail.com

REFERENCES

1. Khalid N, Ahmad SA, Shlofmitz E, Chhabra L. Racial and gender disparities among patients with Takotsubo syndrome. *Clin Cardiol.* 2019;42(1):19.
2. Lemor A, Ramos-Rodriguez AJ, De La Villa R, et al. Impact of gender on in-hospital outcomes in patients with Takotsubo syndrome: a Nationwide analysis from 2006 to 2014. *Clin Cardiol.* 2019;42(1):13-18.
3. Medina de Chazal H, Del Buono MG, Keyser-Marcus L, et al. Stress cardiomyopathy diagnosis and treatment: JACC state-of-the-art review. *J Am Coll Cardiol.* 2018;72(16):1955-1971.